

SPECTRAL DISTORTIONS OF THE COSMIC MICROWAVE BACKGROUND: A NEW WINDOW TO EARLY UNIVERSE PHYSICS

J. Chluba, D. J. Fixsen, M. Kamionkowski, R. Khatri, A. Kogut, J. C. Mather,
M. D. Niemack, J. Silk, R. A. Sunyaev, and E. J. Wollack

Cosmology is now a precise scientific discipline, with detailed theoretical models that fit a wealth of very accurate measurements. Of the many cosmological data sets, the cosmic microwave background (CMB) temperature and polarization *anisotropies* provide the most stringent and robust constraints to theoretical models, allowing us to address fundamental questions about inflation, the nature of dark matter and dark energy, and particle physics [1, 2, 3]. But the CMB holds another, complementary piece of information: its *frequency spectrum*. Since COBE/FIRAS the average CMB spectrum is known to be extremely close to a perfect blackbody, with possible spectral distortions limited to $\Delta I_\nu/I_\nu \lesssim 10^{-5}$ [4, 5]. Although no distortion was detected, this measurement already places tight constraints on the thermal history of our Universe, ruling out cosmologies with extended periods of significant energy release, disturbing the equilibrium between matter and radiation. More than 20 years have past since the launch of COBE and from the technological point of view already today it should be possible to improve the sensitivity by *at least three orders of magnitude* [6]. This opens a new window to the early Universe, on one hand allowing us to directly probe processes that are present within the standard cosmological paradigm, but also opening up a huge discovery space for non-standard physics.

The CMB spectrum constrains energy release occurring at redshift $z \lesssim \text{few} \times 10^6$. A large number of astrophysical or cosmological processes in this range exist, leading to predictions of observable distortions. A small sample of processes follows.

- **REIONIZATION AND STRUCTURE FORMATION:** the first sources of radiation, supernova feedback [7] and structure formation shocks [8, 9, 10] heat the intergalactic medium at low redshifts ($z \lesssim 10$), leading to up-scattering of CMB photons characterized by a Compton y -distortion [11]. The distortion is expected to reach $\Delta I_\nu/I_\nu \simeq 10^{-7} - 10^{-6}$ and thus could be measured at $\simeq 100\sigma$ using present-day technology, teaching us about the average temperature of the intergalactic medium [12], and promising way to find the missing baryons in the local Universe which otherwise are hard to observe [9].
- **INFLATION:** the Silk-damping of small-scale perturbations gives rise to a chemical potential (or μ -distortion) and y -type distortion [13, 14, 15, 16], which directly depends on the shape and amplitude of the primordial power spectrum at scales $0.1 \text{ kpc} \lesssim \lambda \lesssim 1 \text{ Mpc}$ [17, 18]. This allows constraining the trajectory of the inflaton at stages unexplored by CMB anisotropies and other ongoing or planned experiments [19]. The distortion is also sensitive to the difference between adiabatic and isocurvature perturbations [15, 20, 21], as well as primordial non-Gaussianity in the squeezed-limit [22, 23].
- **COSMOLOGICAL RECOMBINATION RADIATION:** the cosmological recombination process of hydrogen and helium introduces distortions at high redshifts ($z \simeq 10^3 - 10^4$), corresponding to $\simeq 260 \text{ kyr}$, $\simeq 130 \text{ kyr}$, and $\simeq 18 \text{ kyr}$ after the big bang. The overall distortions is very small ($\Delta I_\nu/I_\nu \simeq 10^{-9}$) but it has a unique frequency-dependence which opens an independent path to determination of cosmological parameters (like the baryon density and *pre-stellar* helium abundance) and direct measurements of the recombination dynamics, probing the Universe at stages well before the last scattering surface [24].
- **COOLING OF MATTER:** the adiabatic cooling of ordinary matter continuously extracts energy from the CMB photon bath leading to another small but indisputable distortion that directly depends on the baryon density and is characterized by a *negative* μ - and y -parameter [25, 26].

All these examples demonstrate that the CMB spectrum provides a rich and unique source of complementary information about the early Universe, with the certainty for the detection of spectral distortions at a level within reach of present day and future instrumentation. The CMB spectrum could also place interesting constraints on decaying and annihilating particles [27, 28, 29, 25], the power spectrum of small-scale magnetic fields [30], primordial black holes [31], and cosmic strings [32, 33, 34], to mention a few more exotic cases. Deciphering all these signals will be a big challenge for the future, but it holds the potential for new discoveries, providing additional, independent constraints on processes that otherwise will remain a secret of our Universe.

References

- [1] G. F. Smoot, C. L. Bennett, A. Kogut, E. L. Wright, J. Aymon, N. W. Boggess, E. S. Cheng, G. de Amici, S. Gulkis, M. G. Hauser, G. Hinshaw, P. D. Jackson, M. Janssen, E. Kaita, T. Kelsall, and P. Keegstra. Structure in the COBE differential microwave radiometer first-year maps. *ApJL*, 396:L1–L5, September 1992.
- [2] C. L. Bennett, M. Halpern, G. Hinshaw, N. Jarosik, A. Kogut, M. Limon, S. S. Meyer, L. Page, D. N. Spergel, G. S. Tucker, E. Wollack, E. L. Wright, C. Barnes, M. R. Greason, R. S. Hill, E. Komatsu, M. R. Nolte, N. Odegard, H. V. Peiris, and L. Verde. First-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Preliminary Maps and Basic Results. *ApJS*, 148:1–27, September 2003.
- [3] Planck Collaboration, P. A. R. Ade, N. Aghanim, C. Armitage-Caplan, M. Arnaud, M. Ashdown, F. Atrio-Barandela, J. Aumont, C. Baccigalupi, A. J. Banday, and et al. Planck 2013 results. XVI. Cosmological parameters. *ArXiv e-prints*, March 2013.
- [4] J. C. Mather, E. S. Cheng, D. A. Cottingham, R. E. Eplee, Jr., D. J. Fixsen, T. Hewagama, R. B. Isaacman, K. A. Jensen, S. S. Meyer, P. D. Noerdlinger, S. M. Read, and L. P. Rosen. Measurement of the cosmic microwave background spectrum by the COBE FIRAS instrument. *ApJ*, 420:439–444, January 1994.
- [5] D. J. Fixsen, E. S. Cheng, J. M. Gales, J. C. Mather, R. A. Shafer, and E. L. Wright. The Cosmic Microwave Background Spectrum from the Full COBE FIRAS Data Set. *ApJ*, 473:576–+, December 1996.
- [6] A. Kogut, D. J. Fixsen, D. T. Chuss, J. Dotson, E. Dwek, M. Halpern, G. F. Hinshaw, S. M. Meyer, S. H. Moseley, M. D. Seiffert, D. N. Spergel, and E. J. Wollack. The Primordial Inflation Explorer (PIXIE): a nulling polarimeter for cosmic microwave background observations. *JCAP*, 7:25–+, July 2011.
- [7] S. P. Oh, A. Cooray, and M. Kamionkowski. Sunyaev-Zeldovich fluctuations from the first stars? *MNRAS*, 342:L20–L24, June 2003.
- [8] R. A. Sunyaev and Y. B. Zeldovich. Formation of Clusters of Galaxies; Protocluster Fragmentation and Intergalactic Gas Heating. *A&A*, 20:189–+, August 1972.
- [9] R. Cen and J. P. Ostriker. Where Are the Baryons? *ApJ*, 514:1–6, March 1999.
- [10] F. Miniati, D. Ryu, H. Kang, T. W. Jones, R. Cen, and J. P. Ostriker. Properties of Cosmic Shock Waves in Large-Scale Structure Formation. *ApJ*, 542:608–621, October 2000.
- [11] Y. B. Zeldovich and R. A. Sunyaev. The Interaction of Matter and Radiation in a Hot-Model Universe. *Ap&SS*, 4:301–316, July 1969.
- [12] P. Zhang, U.-L. Pen, and H. Trac. The temperature of the intergalactic medium and the Compton y parameter. *MNRAS*, 355:451–460, December 2004.
- [13] R. A. Sunyaev and Y. B. Zeldovich. Small scale entropy and adiabatic density perturbations – Antimatter in the Universe. *Ap&SS*, 9:368–382, December 1970.
- [14] R. A. Daly. Spectral distortions of the microwave background radiation resulting from the damping of pressure waves. *ApJ*, 371:14–28, April 1991.
- [15] J. D. Barrow and P. Coles. Primordial density fluctuations and the microwave background spectrum. *MNRAS*, 248:52–57, January 1991.
- [16] W. Hu, D. Scott, and J. Silk. Power spectrum constraints from spectral distortions in the cosmic microwave background. *ApJL*, 430:L5–L8, July 1994.
- [17] J. Chluba, R. Khatri, and R. A. Sunyaev. CMB at 2×2 order: the dissipation of primordial acoustic waves and the observable part of the associated energy release. *MNRAS*, 425:1129–1169, September 2012.
- [18] R. Khatri, R. A. Sunyaev, and J. Chluba. Mixing of blackbodies: entropy production and dissipation of sound waves in the early Universe. *A&A*, 543:A136, July 2012.
- [19] J. Chluba, A. L. Erickcek, and I. Ben-Dayan. Probing the Inflaton: Small-scale Power Spectrum Constraints from Measurements of the Cosmic Microwave Background Energy Spectrum. *ApJ*, 758:76, October 2012.
- [20] W. Hu and N. Sugiyama. Thermal history constraints on the isocurvature baryon model. *ApJ*, 436:456–466, December 1994.
- [21] J. B. Dent, D. A. Easson, and H. Tashiro. Cosmological constraints from CMB distortion. *Phys.Rev.D*, 86(2):023514, July 2012.
- [22] E. Pajer and M. Zaldarriaga. New Window on Primordial Non-Gaussianity. *Physical Review Letters*, 109(2):021302, July 2012.
- [23] J. Ganc and E. Komatsu. Scale-dependent bias of galaxies and μ -type distortion of the cosmic microwave background spectrum from single-field inflation with a modified initial state. *Phys.Rev.D*, 86(2):023518, July 2012.
- [24] R. A. Sunyaev and J. Chluba. Signals from the epoch of cosmological recombination (Karl Schwarzschild Award Lecture 2008). *Astronomische Nachrichten*, 330:657–+, 2009.
- [25] J. Chluba and R. A. Sunyaev. The evolution of CMB spectral distortions in the early Universe. *MNRAS*, 419:1294–1314, January 2012.
- [26] R. Khatri, R. A. Sunyaev, and J. Chluba. Does Bose-Einstein condensation of CMB photons cancel μ distortions created by dissipation of sound waves in the early Universe? *arXiv:1110.0475*, October 2011.
- [27] W. Hu and J. Silk. Thermalization constraints and spectral distortions for massive unstable relic particles. *Physical Review Letters*, 70:2661–2664, May 1993.

- [28] P. McDonald, R. J. Scherrer, and T. P. Walker. Cosmic microwave background constraint on residual annihilations of relic particles. *Phys.Rev.D*, 63(2):023001—+, January 2001.
- [29] J. Chluba. Could the cosmological recombination spectrum help us understand annihilating dark matter? *MNRAS*, 402:1195–1207, February 2010.
- [30] K. Jedamzik, V. Katalinić, and A. V. Olinto. Limit on Primordial Small-Scale Magnetic Fields from Cosmic Microwave Background Distortions. *Physical Review Letters*, 85:700–703, July 2000.
- [31] B. J. Carr, K. Kohri, Y. Sendouda, and J. Yokoyama. New cosmological constraints on primordial black holes. *Phys.Rev.D*, 81(10):104019—+, May 2010.
- [32] J. P. Ostriker and C. Thompson. Distortion of the cosmic background radiation by superconducting strings. *ApJL*, 323:L97–L101, December 1987.
- [33] H. Tashiro, E. Sabancilar, and T. Vachaspati. CMB distortions from superconducting cosmic strings. *Phys.Rev.D*, 85(10):103522, May 2012.
- [34] H. Tashiro, E. Sabancilar, and T. Vachaspati. CMB Distortions from Damping of Acoustic Waves Produced by Cosmic Strings. *ArXiv:1212.3283*, December 2012.